326 U.S. A THE ARCHENERGY

RC COMMISSION

Collection SEC SEARONG

Box 69

OHM INCIDENCE OF THYRIOD

FOIDER OR OTHER DISEASES

N THE EXPOSED GR

UNITED STATES
ATOMIC ENERGY COMMISSION
WASHINGTON, D.C. 20545

403534

DC FILE

January 6, 1965

JSN 1 (8/65

MEMORANDUM FOR CHAIRMAN SEABORG

1965_1968 COMMISSIONER BUNTING

COMMISSIONER PALFREY
COMMISSIONER RAMEY
COMMISSIONER TAPE

THROUGH GENERAL MANAGE

fak V. Vinaguerra JAN 7 196

SUBJECT: FURTHER INFORMATION ON THREE MARSHALLESE GIRLS OPERATED ON FOR REMOVAL OF THYROID NODULES (SEE ATTACHED COPY OF PREVIOUS MEMO, THIS SUBJECT).

Dr. Robert A. Conard, Medical Department, Brookhaven National Laboratory, who has responsibility for conducting the clinical surveys, reports: "Sections of the nodules have been examined by a number of pathologists including Drs. S. Warren, G. H. Klink, H. A. Johnson and S. Lindsay. The concensus is that the nodules are benign, though with certain bizarre areas, and resemble iodine deficiency adenomata."

Dr. Lindsay noted that the nodules in the Rongelap cases are similar to ones he has seen in children following I¹³¹ therapy. This Division has a file of Dr. Lindsay's histological reports and a review paper by Lindsay and Chaikoff, <u>Cancer Research</u>, 24:1099, 1964.

The radiation dosages which these children's thyroids may have received are being re-evaluated by Dr. John Gofman's group, Lawrence Radiation Laboratory.

It should be added that nodules in the thyroid glands of young people frequently become malignant in the course of time. Therefore, while these which were removed promptly after discovery were not malignant, there is a chance that in time they might have assumed neoplastic characteristics. Dr. Conard next clinical survey will take place in February 1965.

C. L. Dunham, M.D.

Director

Division of Biology and Medicine

Attachment:

Copy of memo dated 10/2/64 from C. L. Dunham to Chairman Seaborg